


PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 471/04, 487/04, A61K 31/505	A1	(11) International Publication Number: WO 98/23613 (43) International Publication Date: 4 June 1998 (04.06.98)
(21) International Application Number: PCT/IB97/01393 (22) International Filing Date: 5 November 1997 (05.11.97) (30) Priority Data: 60/031,862 27 November 1996 (27.11.96) US 60/041,846 9 April 1997 (09.04.97) US (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ARNOLD, Lee, Daniel [CA/US]; 216 Ruggles Street, Westborough, MA 01581 (US). MOYER, Mikel, Paul [US/US]; 5 Butterwick Lane, Old Lyme, CT 06371 (US). SOBOLOV-JAYNES, Susan, Beth [US/US]; 4 Piney Branch Road, Ivoryton, CT 06442 (US). (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: FUSED BICYCLIC PYRIMIDINE DERIVATIVES <div style="text-align: center;"> (I)</div> (57) Abstract <p>The invention relates to compounds of formula (I) and to pharmaceutically acceptable salts thereof, wherein R¹, R² and Z are as defined herein. The invention also relates to pharmaceutical compositions containing the compounds of formula (I) and to methods of using said compounds in the treatment of hyperproliferative diseases such as cancer.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

FUSED BICYCLIC PYRIMIDINE DERIVATIVESBackground of the Invention

This invention relates to novel bicyclic pyrimidine derivatives that are useful in the treatment of hyperproliferative diseases, such as cancers, in mammals. This invention also relates to a method of using such compounds in the treatment of
10 hyperproliferative diseases in mammals, especially humans, and to pharmaceutical compositions containing such compounds.

Compounds that are useful in the treatment of hyperproliferative diseases are also disclosed in the following co-pending patent applications: PCT international patent application number PCT/IB97/00675 (filed June 11, 1997), United States provisional
15 patent application number 60/028881 (filed October 17, 1996), PCT international patent application number PCT/IB97/00584 (filed May 22, 1997), United States patent application number 08/653,786 (filed May 28, 1996), PCT international patent application publication number WO 96/40142 (published December 19, 1996), and PCT international patent application publication number WO 95/23141 (published August 31,
20 1995). Each of the foregoing United States and PCT international patent applications is incorporated herein by reference in its entirety.

It is known that a cell may become cancerous by virtue of the transformation of a portion of its DNA into an oncogene (i.e. a gene that upon activation leads to the formation of malignant tumor cells). Many oncogenes encode proteins which are
25 aberrant tyrosine kinases capable of causing cell transformation. Alternatively, the overexpression of a normal proto-oncogenic tyrosine kinase may also result in proliferative disorders, sometimes resulting in a malignant phenotype.

Receptor tyrosine kinases are large enzymes that span the cell membrane and possess an extracellular binding domain for growth factors such as epidermal growth
30 factor, a transmembrane domain, and an intracellular portion that functions as a kinase to phosphorylate specific tyrosine residue in proteins and hence to influence cell proliferation. It is known that such kinases are often aberrantly expressed in common human cancers such as breast cancer, gastrointestinal cancer such as colon, rectal or stomach cancer, leukemia, and ovarian, bronchial or pancreatic cancer. It has also
35 been shown that epidermal growth factor receptor (EGFR), which possesses tyrosine kinase activity, is mutated or overexpressed in many human cancers such as brain, lung, squamous cell, bladder, gastric, breast, head and neck, oesophageal, gynecological and thyroid cancers. Thus, it is believed that inhibitors of receptor

-2-


tyrosine kinases, such as the compounds of the present invention, are useful as selective inhibitors of the growth of mammalian cancer cells.

It has also been shown that EGFR inhibitors may be useful in the treatment of pancreatitis and kidney disease (such as proliferative glomerulonephritis and diabetes-induced renal disease), and may reduce successful blastocyte implantation and therefore may be useful as a contraceptive. See PCT international application publication number WO 95/19970 (published July 27, 1995).

It is known that polypeptide growth factors such as vascular endothelial growth factor (VEGF) having a high affinity to the human kinase insert-domain-containing receptor (KDR) or the murine fetal liver kinase 1 (FLK-1) receptor have been associated with the proliferation of endothelial cells and more particularly vasculogenesis and angiogenesis. See PCT international application publication number WO 95/21613 (published August 17, 1995). Agents, such as the compounds of the present invention, that are capable of binding to or modulating the KDR/FLK-1 receptor may be used to treat disorders related to vasculogenesis or angiogenesis such as diabetes, diabetic retinopathy, hemangioma, glioma, melanoma, Kaposi's sarcoma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer.

Summary of the Invention


The present invention relates to compounds of the formula



and to pharmaceutically acceptable salts thereof, wherein:

-3-

Z is a group of the formula




wherein n is an integer from 0 to 2 and p is an integer from 0 to 3;

R¹ is H, C₁-C₆ alkyl or -C(O)(C₁-C₆ alkyl);


-4-

R^2 is phenyl or 1H-indazol-5-yl, wherein said groups are optionally substituted by 1 to 3 R^5 substituents, or R^2 is a group of the formula (Ii) or (Ij)



(Ii)


or



(Ij)

wherein p is an integer from 0 to 3 and n is an integer from 0 to 2;

or R^1 and R^2 are taken together to form a group of the formula (Ik)



20

;

(Ik)

25

wherein the dashed line indicates a single or double bond and m is an integer from 0 to 4;

each R^3 is independently H, $-C(O)OR^9$, or C_1-C_6 alkyl wherein said alkyl is optionally substituted by halo, $-OR^9$, $-NR^9R^{10}$, or $-C(O)OR^9$;

30 R^4 is R^3 , $-OR^9$, or $-NR^9R^{10}$;

each R^5 is independently halo, cyano, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, $-OR^9$, $-NR^9R^{10}$, nitro, or C_6-C_{10} aryl wherein said alkyl, alkenyl, alkynyl and aryl R^5

-5-

groups are optionally substituted by 1 to 3 substituents independently selected from halo, nitro, C₁-C₄ alkyl and -OR⁹;

R⁶ and R⁷ are independently H or R⁵;

R⁸ is H, -SO₂(C₆-C₁₀ aryl), -(CH₂)_q(5-10 membered heterocyclyl), C₂-C₆ alkenyl,
 5 C₁-C₆ alkyl, -(CH₂)_qO(CH₂)_q(C₁-C₆ alkoxy), -(CH₂)_q(C₁-C₆ alkoxy), -C(O)(C₁-C₆ alkoxy),
 or -SO₂(C₁-C₄ alkyl) wherein each q is independently an integer from 2 to 4;

each R⁹ and R¹⁰ is independently H or C₁-C₆ alkyl; and,

R¹¹ is trifluoromethyl, halo, nitro, -OR⁹, -NR⁹R¹⁰, cyano, C₁-C₄ alkyl, -S(O)_xR⁹
 wherein x is an integer from 0 to 2, -C(O)OR⁹, -OC(O)(C₁-C₄ alkyl), -C(O)NR⁹R¹⁰,
 10 -NR⁹C(O)(C₁-C₄ alkyl), -C(O)NHSO₂(C₁-C₄ alkyl), -NHCH₂C(O)NR⁹R¹⁰, -NHC(O)(C₁-C₄
 alkoxy), -NHOC(O)(C₁-C₄ alkyl), -NR⁹OR¹⁰, anilino, pyrrolidinyl, piperidinyl, azido,
 guanidino, morpholino, phenyl, -C(O)(C₁-C₆ alkyl), benzenesulfonyl, allyl, thiophenyl,
 piperazinyl, 4-(C₁-C₄ alkyl)-piperazinyl, phenylthio, benzenesulphonamido, 2-
 oxopyrrolidin-1-yl, 2,5-dioxopyrrolidin-1-yl, phenoxy, benzoyloxy, benzoylamino,
 15 -(CH₂)_wO(CH₂)_vOR⁹, -O(CH₂)_wO(CH₂)_vOR⁹, -O(CH₂)_wC(O)OR⁹, -O(CH₂)_wC(O)NR⁹R¹⁰,
 -(CH₂)_wS(CH₂)_vOR⁹, -NH(CH₂)_vO(C₁-C₄ alkyl), -NH(CH₂)_w(C₆-C₁₀ aryl),
 -NHC(O)(CH₂)_w(C₁-C₄ alkoxy), or -O(CH₂)_w(C₆-C₁₀ aryl), wherein w is an integer from 1
 to 4 and v is an integer from 2 to 4, and wherein the alkyl, heterocyclic, and aryl
 moieties of the foregoing R¹¹ groups are optionally substituted by 1 or 2 substituents
 20 independently selected from the group consisting of halo, C₁-C₄ alkyl, -OR⁹, -NR⁹R¹⁰,
 -C(O)OR⁹, -OC(O)(C₁-C₄ alkyl), -C(O)NR⁹R¹⁰, -NHC(O)(C₁-C₄ alkyl), nitro, imidazolyl,
 piperidino, morpholino, and piperazinyl;

with the proviso that where Z is a group of the formula (Ie), and R² is phenyl,
 then said phenyl is substituted by 1 to 3 substituents independently selected from C₁-C₆
 25 alkyl, C₂-C₆ alkynyl and halo, and one of R⁶ and R⁷ is halo or H and the other is as
 defined above; and,

with the further proviso that where Z is a group of the formula (Ia), (Ib), (Ic) or
 (Id), and R² is phenyl, then said phenyl is substituted by C₂-C₆ alkynyl.

Preferred compounds of formula I include those wherein R² is optionally
 30 substituted phenyl. More preferred are those compounds of formula I wherein R² is
 phenyl substituted by C₂-C₆ alkynyl, in particular ethynyl.

Other preferred compounds of formula I include those wherein Z is a pyrrolo
 moiety of formula (Ie) or (If).

Other preferred compounds of formula I include those wherein R¹ and R² are taken together to form an indole or indoline moiety of formula (Ik).

Specific preferred compounds of formula I include the following:

- 4-(6-Chloro-2,3-dihydro-indol-1-yl)-7H-pyrrolo[2,3-d]pyrimidine;
- 5 4-(6-Methyl-2,3-dihydro-indol-1-yl)-7H-pyrrolo[2,3-d]pyrimidine;
- 4-(6-Chloro-5-fluoro-2,3-dihydro-indol-1-yl)-7H-pyrrolo[2,3-d]pyrimidine;
- 1-(4-m-Tolylamino-pyrrolo[2,3-d]pyrimidin-7-yl)-ethanone;
- 4-(6-Chloro-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- 4-(6-Bromo-5-chloro-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- 10 4-(6-Fluoro-5-chloro-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- 4-(6-Iodo-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- (7-Benzenesulfonyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-(3-ethynyl-phenyl)-amine;
- 4-(6-Chloro-2,3-dihydro-indol-1-yl)-5H-pyrrolo[3,2-d]pyrimidin-6-ol;
- (3-Ethynyl-phenyl)-[7-(2-morpholin-4-yl-ethyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-
- 15 amine;
- (3-Ethynyl-phenyl)-[7-(2-methoxy-ethyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine;
- (3-Ethynyl-phenyl)-{7-[2-(2-methoxy-ethoxy)-ethyl]-7H-pyrrolo[2,3-d]pyrimidin-4-yl}-amine;
- (7-Allyl-pyrrolo[2,3-d]pyrimidin-4-yl)-(3-ethynyl-phenyl)-amine;
- 20 N-(5-Iodo-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-N-m-tolyl-acetamide;
- 4-(6-Chloro-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6-Bromo-5-fluoro-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6-Chloro-5-fluoro-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6-Iodo-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 25 4-(4-Bromo-7-methyl-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6-Bromo-7-methyl-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6,7-Dimethyl-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- (3-Ethynyl-phenyl)-pyrido[3,4-d]pyrimidin-4-yl-amine;
- Benzo[b]thiophen-5-yl-pyrido[3,4-d]pyrimidin-4-yl-amine;
- 30 (3-Ethynyl-phenyl)-(5-p-tolyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;
- (3-Ethynyl-phenyl)-(5-thiophen-2-yl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;
- (3-Ethynyl-phenyl)-[5-(4-methoxy-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine;
- (3-Ethynyl-phenyl)-[5-(3-nitro-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine;

-7-

[5-(4-Chloro-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-(3-ethynyl-phenyl)-amine;
(3-Bromo-phenyl)-(6-bromo-5-phenyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;
and the pharmaceutically acceptable salts of the foregoing compounds.

The invention also relates to a pharmaceutical composition for the treatment of
5 a hyperproliferative disorder in a mammal which comprises a therapeutically effective
amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, and
a pharmaceutically acceptable carrier. In one embodiment, said pharmaceutical
composition is for the treatment of cancer such as brain, lung, squamous cell, bladder,
gastric, pancreatic, breast, head, neck, renal, kidney, ovarian, prostate, colorectal,
10 oesophageal, gynecological or thyroid cancer. In another embodiment, said
pharmaceutical composition is for the treatment of a non-cancerous hyperproliferative
disorder such as benign hyperplasia of the skin (e.g., psoriasis) or prostate (e.g.,
benign prostatic hypertrophy (BPH)).

The invention also relates to a pharmaceutical composition for the treatment of
15 a hyperproliferative disorder in a mammal which comprises a therapeutically effective
amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, in
combination with an anti-tumor agent selected from the group consisting of mitotic
inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, enzymes,
topoisomerase inhibitors, biological response modifiers, anti-hormones, and anti-
20 androgens, and a pharmaceutically acceptable carrier.

The invention also relates to a pharmaceutical composition for the treatment of
pancreatitis or kidney disease (including proliferative glomerulonephritis and diabetes-
induced renal disease) in a mammal which comprises a therapeutically effective
amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, and
25 a pharmaceutically acceptable carrier.

The invention also relates to a pharmaceutical composition for the prevention
of blastocyte implantation in a mammal which comprises a therapeutically effective
amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, and
a pharmaceutically acceptable carrier.

30 The invention also relates to a pharmaceutical composition for treating a disease
related to vasculogenesis or angiogenesis in a mammal which comprises a
therapeutically effective amount of a compound of formula I, or a pharmaceutically
acceptable salt thereof, and a pharmaceutically acceptable carrier. In one embodiment,

-8-

said pharmaceutical composition is for treating a disease selected from the group consisting of diabetes, diabetic retinopathy, hemangioma, glioma, melanoma, Kaposi's sarcoma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer.

The invention also relates to a method of treating a hyperproliferative disorder
5 in a mammal which comprises administering to said mammal a therapeutically effective amount of the compound of formula I or a pharmaceutically acceptable salt thereof. In one embodiment, said method relates to the treatment of cancer such as brain, squamous cell, bladder, gastric, pancreatic, breast, head, neck, oesophageal, prostate, colorectal, lung, renal, kidney, ovarian, gynecological or thyroid cancer. In another
10 embodiment, said method relates to the treatment of a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (e.g., psoriasis) or prostate (e.g., benign prostatic hypertrophy (BPH)).

The invention also relates to a method for the treatment of a hyperproliferative disorder in a mammal which comprises administering to said mammal a therapeutically
15 effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with an anti-tumor agent selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, and anti-androgens.

20 The invention also relates to a method of treating pancreatitis or kidney disease in a mammal which comprises administering to said mammal a therapeutically effective amount of the compound of formula I or a pharmaceutically acceptable salt thereof.

The invention also relates to a method of preventing blastocyte implantation in a mammal which comprises administering to said mammal a therapeutically effective
25 amount of the compound of formula I or a pharmaceutically acceptable salt thereof.

The invention also relates to a method of treating diseases related to vasculogenesis or angiogenesis in a mammal which comprises administering to said mammal an effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof. In one embodiment, said method is for treating a disease
30 selected from the group consisting of diabetes, diabetic retinopathy, hemangioma, glioma, melanoma, Kaposi's sarcoma and ovarian, breast, lung, pancreatic, prostate, colon and epidermoid cancer.

Patients that can be treated with compounds of the formula I, and the pharmaceutically acceptable salts of said compounds, according to the methods of this invention include, for example, patients that have been diagnosed as having psoriasis, BPH, lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head and neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer, gynecologic tumors (e.g., uterine sarcomas, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina or carcinoma of the vulva), Hodgkin's disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system (e.g., cancer of the thyroid, parathyroid or adrenal glands), sarcomas of soft tissues, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute leukemia, solid tumors of childhood, lymphocytic lymphomas, cancer of the bladder, cancer of the kidney or ureter (e.g., renal cell carcinoma, carcinoma of the renal pelvis), or neoplasms of the central nervous system (e.g., primary CNS lymphoma, spinal axis tumors, brain stem gliomas or pituitary adenomas).

The term "halo", as used herein, unless otherwise indicated, means fluoro, chloro, bromo or iodo. Preferred halo groups are fluoro, chloro and bromo.

The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, cyclic or branched moieties. It is understood that for cyclic moieties at least three carbon atoms are required in said alkyl group.

The term "alkoxy", as used herein, unless otherwise indicated, includes O-alkyl groups wherein "alkyl" is as defined above.

The term "aryl", as used herein, unless otherwise indicated, includes an organic radical derived from an aromatic hydrocarbon by removal of one hydrogen, such as phenyl or naphthyl.

The term "5-10 membered heterocyclyl", as used herein, unless otherwise indicated, includes aromatic and non-aromatic heterocyclic groups containing one or more heteroatoms each selected from O, S and N, wherein each heterocyclic group has from 5-10 atoms in its ring system. The heterocyclic groups include benzo-fused ring systems and ring systems substituted with one or more oxo moieties. An example of a 5 membered heterocyclic group is thiazolyl, and an example of a 10 membered heterocyclic group is quinolinyl. Examples of non-aromatic heterocyclic groups are

pyrrolidinyl, piperidino, morpholino, thiomorpholino and piperazinyl. Examples of aromatic heterocyclic groups are pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl and thiazolyl. Heterocyclic groups having a fused benzene ring include benzimidazolyl.

5 The phrase "pharmaceutically acceptable salt(s)", as used herein, unless otherwise indicated, includes salts of acidic or basic groups which may be present in the compounds of formula I. The compounds of formula I that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of
10 such basic compounds of formula I are those that form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, acid citrate, tartrate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucaronate,
15 saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts.


Those compounds of the formula I that are acidic in nature, are capable of forming base salts with various pharmacologically acceptable cations. Examples of
20 such salts include the alkali metal or alkaline earth metal salts and particularly, the sodium and potassium salts.

Certain compounds of formula I may have asymmetric centers and therefore exist in different enantiomeric forms. This invention relates to the use of all optical isomers and stereoisomers of the compounds of formula I and mixtures thereof. The
25 compounds of formula I may also exist as tautomers. This invention relates to the use of all such tautomers and mixtures thereof.


Detailed Description of the Invention

The preparation of the compounds of the present invention is illustrated in the following Schemes 1 and 2.

-11-

Scheme 1

-12-

Scheme 2

IV



V

25

-13-

The compounds of the present invention are readily prepared according to synthetic methods familiar to those skilled in the art. Such methods are disclosed in PCT international patent application publication number WO 96/40142 (published December 19, 1996), referred to above, and in PCT international application publication
5 numbers WO 95/19774 (published July 27, 1995) and WO 95/19970 (published July 27, 1995), each of which is incorporated herein by reference.

Scheme 1 illustrates the coupling of the bicyclic compound of formula II with the amine of formula III to provide the compound of formula I. In the compounds of formulas II and III, X is hydroxy or chloro and Z, R¹ and R² are as defined above. In
10 general, the compound of formula II is coupled with the amine of formula III in a solvent, such as a C₁-C₆ alcohol, dimethylformamide (DMF), N-methylpyrrolidin-2-one (NMP), chloroform, acetonitrile, tetrahydrofuran (THF), dimethylsulfoxide (DMSO), 1,4-dioxane or pyridine, optionally in the presence of a base, such as pyridine or triethylamine, and optionally in the presence of pyridine hydrochloride as a catalyst,
15 under an inert atmosphere, such as dry nitrogen gas, at a temperature of from ambient to reflux temperature, preferably 80-125°C, for a period of about 2 hours to 72 hours.

The bicyclic compound of formula II is prepared according to synthetic methods known to those skilled in the art. Such methods are disclosed in WO 95/19774 and WO 95/19970, referred to above. Other methods of preparing the compounds of
20 formula II are also disclosed in PCT international application publication number WO 92/12718 (published August 6, 1992), A. Petric et al., *Fur Chemie* 114, 615-624 (1983) and *Nucleic Acids Research*, v. 12, no. 2 (1984). Where the compound of formula III is an optionally substituted indole or indoline moiety, such compounds can be prepared according to one or more methods known to those skilled in the art. Such methods are
25 described in PCT international patent application publication number WO 95/23141, referred to above, and in W.C. Sumpter and F.M. Miller, "Heterocyclic Compounds with Indole and Carbazole Systems," in volume 8 of "The Chemistry of Heterocyclic Compounds", Interscience Publishers Inc., New York (1954). Optional substituents can be included as appropriate before or after the coupling step illustrated in Scheme 1.
30 Prior to the coupling step, primary and secondary amino moieties (other than the amine of formula III) are preferably protected using a nitrogen protecting group known to those skilled in the art. Such protecting groups and their use are described in T.W. Greene

and P.G.M. Wuts, "Protective Groups in Organic Synthesis," Second Edition, John Wiley & Sons, New York, 1991.

Scheme 2 illustrates the cyclization of the compound of formula IV (wherein Me means methyl) to provide the compound of formula V. This illustrates the introduction of the Z moiety in the compound of formula I after the amino ($-NR^1R^2$) moiety has been coupled with the pyrimidine ring. While Scheme 2 illustrates the formation of a pyrrolo[3,2-d]pyrimidine ring system, other ring systems can be made by one or more analogous methods disclosed in WO 90/19774, referred to above. In Scheme 2, the compound of formula IV, in an alcoholic solvent, such as ethanol, is catalytically hydrogenated using an appropriate catalyst, such as 10% palladium on carbon, under an H_2 atmosphere at ambient temperature for a period of about three hours to provide the compound of formula V. Optional substituents can be introduced into the compound of formula V, as appropriate, according to methods known to those skilled in the art, to provide the desired compound of formula I.

The compounds of the present invention may have asymmetric carbon atoms. Such diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known to those skilled in the art, for example, by chromatography or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixtures into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., alcohol), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. All such isomers, including diastereomer mixtures and pure enantiomers are considered as part of the invention.

The compounds of formula I that are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate the compound of formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent,

-15-

such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained. The desired acid salt can also be precipitated from a solution of the free base in an organic solvent by adding to the solution an appropriate mineral or organic acid.

5 Those compounds of the formula I that are acidic in nature, are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically
10 acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of formula I. Such non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium calcium and magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically
15 acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed
20 in order to ensure completeness of reaction and maximum yields of the desired final product.

 The active compounds of this invention are potent inhibitors of the *erbB* family of oncogenic and protooncogenic protein tyrosine kinases such as epidermal growth factor receptor (EGFR), *erbB2*, *HER3*, or *HER4* and thus are all adapted to therapeutic
25 use as antiproliferative agents (e.g., anticancer) in mammals, particularly in humans. In particular, the compounds of this invention are useful in the prevention and treatment of a variety of human hyperproliferative disorders such as malignant and benign tumors of the liver, kidney, bladder, breast, gastric, ovarian, colorectal, prostate, pancreatic, lung, vulval, thyroid, hepatic carcinomas, sarcomas, glioblastomas, head and neck, and
30 other hyperplastic conditions such as benign hyperplasia of the skin (e.g., psoriasis) and benign hyperplasia of the prostate (e.g., BPH). It is, in addition, expected that a compound of the present invention may possess activity against a range of leukemias and lymphoid malignancies.

-16-

The active compounds may also be useful in the treatment of additional disorders in which aberrant expression ligand/receptor interactions or activation or signalling events related to various protein tyrosine kinases, are involved. Such disorders may include those of neuronal, glial, astrocytal, hypothalamic, and other glandular, macrophagal, epithelial, stromal, and blastocoelic nature in which aberrant function, expression, activation or signalling of the erbB tyrosine kinases are involved. In addition, compounds of formula I may have therapeutic utility in inflammatory, angiogenic and immunologic disorders involving both identified and as yet unidentified tyrosine kinases that are inhibited by compounds of the formula I.

10 The in vitro activity of the active compounds in inhibiting the receptor tyrosine kinase (and thus subsequent proliferative response, e.g., cancer) may be determined by the following procedure.

Activity of the active compounds, *in vitro*, can be determined by the amount of inhibition of the phosphorylation of an exogenous substrate (e.g., Lys₃ - Gastrin or polyGluTyr (4:1) random copolymer (I. Posner et al., J. Biol. Chem. 267 (29), 20638-47 (1992)) on tyrosine by epidermal growth factor receptor kinase by a test compound relative to a control. Affinity purified, soluble human EGF receptor (96 ng) is obtained according to the procedure in G. N. Gill, W. Weber, Methods in Enzymology 146, 82-88 (1987) from A431 cells (American Type Culture Collection, Rockville, MD) and preincubated in a microfuge tube with EGF (2µg/ml) in phosphorylation buffer + vanadate (PBV: 50 mM HEPES, pH 7.4; 125 mM NaCl; 24 mM MgCl₂; 100 µM sodium orthovanadate), in a total volume of 10 µl, for 20-30 minutes at room temperature. The test compound, dissolved in dimethylsulfoxide (DMSO), is diluted in PBV, and 10 µl is mixed with the EGF receptor /EGF mix, and incubated for 10-30 minutes at 30°C. The phosphorylation reaction is initiated by addition of 20 µl ³³P-ATP/ substrate mix (120 µM Lys₃-Gastrin (sequence in single letter code for amino acids, KKKGPWLEEEEEAYGWLDF), 50 mM Hepes pH 7.4, 40 µM ATP, 2 µCi γ-[³³P]-ATP) to the EGFr/EGF mix and incubated for 20 minutes at room temperature. The reaction is stopped by addition of 10 µl stop solution (0.5 M EDTA, pH 8; 2mM ATP) and 6 µl 2N HCl. The tubes are centrifuged at 14,000 RPM, 4°C, for 10 minutes. 35 µl of supernatant from each tube is pipetted onto a 2.5 cm circle of Whatman P81 paper, bulk washed four times in 5% acetic acid, 1 liter per wash, and then air dried. This results in the binding of substrate to the paper with loss of free ATP on washing. The

-17-

[³³P] incorporated is measured by liquid scintillation counting. Incorporation in the absence of substrate (e.g., lys₃-gastrin) is subtracted from all values as a background and percent inhibition is calculated relative to controls without test compound present.

Such assays, carried out with a range of doses of test compounds, allow the
5 determination of an approximate IC₅₀ value for the *in vitro* inhibition of EGFR kinase activity. The compounds of the formula I that were tested using the procedure described above exhibited IC₅₀ values in the range of 0.0001-30 μM.

Activity of the active compounds, *in vivo*, can be determined by the amount of inhibition of tumor growth by a test compound relative to a control. The tumor growth
10 inhibitory effects of various compounds are measured according to the methods of Corbett T. H., et al. "Tumor Induction Relationships in Development of Transplantable Cancers of the Colon in Mice for Chemotherapy Assays, with a Note on Carcinogen Structure", Cancer Res., 35, 2434-2439 (1975) and Corbett, T. H., et al., "A Mouse Colon-tumor Model for Experimental Therapy", Cancer Chemother. Rep. (Part 2)", 5,
15 169-186 (1975), with slight modifications. Tumors are induced in the left flank by s.c. injection of 1 X 10⁶ log phase cultured tumor cells (human MDA-MB-468 breast or human HN5 head and neck carcinoma cells) suspended in 0.10 ml RPMI 1640. After sufficient time has elapsed for the tumors to become palpable (2-3 mm in diameter) the test animals (athymic mice) are treated with active compound (formulated by dissolution
20 in DMSO typically at a concentration of 50 to 100 mg/mL followed by 1:9 dilution into saline or, alternatively, 1:9 dilution into 0.1% Pluronic[®] P105 in 0.9% saline) by the intraperitoneal (ip) or oral (po) routes of administration twice daily (i.e., every 12 hours) for 5 consecutive days. In order to determine an anti-tumor effect, the tumor is measured in millimeters with Vernier calipers across two diameters and the tumor size
25 (mg) is calculated using the formula: Tumor weight = (length x [width]²)/2, according to the methods of Geran, R.I., et al. "Protocols for Screening Chemical Agents and Natural Products Against Animal Tumors and Other Biological Systems", Third Edition, Cancer Chemother. Rep., 3, 1-104 (1972). Results are expressed as percent inhibition, according to the formula: Inhibition (%) = (TuW_{control} - TuW_{test})/TuW_{control} x 100%. The
30 flank site of tumor implantation provides reproducible dose/response effects for a variety of chemotherapeutic agents, and the method of measurement (tumor diameter) is a reliable method for assessing tumor growth rates. The title compounds of the

-18-

experimental examples of this case that are compounds of the formula I all exhibited, when tested in the above assay, percent inhibition values greater than 50% at 10 μ M.

Other methods of assessing the activity of the compounds of the present invention are referred to in PCT international application publication number WO
5 95/21613 (published August 17, 1995) which incorporated herein by reference.

Administration of the active compounds can be effected by any method that enables delivery of the compounds to the site of action (e.g., cancer cells). These methods include oral routes, intraduodenal routes, parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion), topical
10 administration, etc.

The amount of the active compound administered will be dependent on the subject being treated, the severity of the disorder or condition, the rate of administration and the judgement of the prescribing physician. However, an effective dosage is in the range of about 0.001 to about 100 mg per kg body weight per day, preferably about 1
15 to about 35 mg/kg/day, in single or divided doses. For a 70 kg human, this would amount to about 0.05 to about 7 g/day, preferably about 0.2 to about 2.5 g/day. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into
20 several small doses for administration throughout the day.

The active compound may be applied as a sole therapy or may involve one or more other anti-tumour substances, for example those selected from, for example, mitotic inhibitors, for example vinblastine; alkylating agents, for example cis-platin, carboplatin and cyclophosphamide; anti-metabolites, for example 5-fluorouracil,
25 cytosine arabinoside and hydroxyurea, or, for example, one of the preferred anti-metabolites disclosed in European Patent Application No. 239362 such as N-(5-[N-(3,4-dihydro-2-methyl-4-oxoquinazolin-6-ylmethyl)-N-methylamino]-2-thenoyl)-L-glutamic acid; intercalating antibiotics, for example adriamycin and bleomycin; enzymes, for example interferon; and anti-hormones, for example anti-estrogens such as NolvadexTM
30 (tamoxifen) or, for example anti-androgens such as CasodexTM (4'-cyano-3-(4-fluorophenylsulphonyl)-2-hydroxy-2-methyl-3'-(trifluoromethyl)propionanilide). Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate dosing of the individual components of the treatment.

The pharmaceutical composition may, for example, be in a form suitable for oral administration as a tablet, capsule, pill, powder, sustained release formulations, solution, suspension, for parenteral injection as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository. The pharmaceutical composition may be in unit dosage forms suitable for single administration of precise dosages. The pharmaceutical composition will include a conventional pharmaceutical carrier or excipient and a compound according to the invention as an active ingredient. In addition, it may include other medicinal or pharmaceutical agents, carriers, adjuvants, etc.

10 Exemplary parenteral administration forms include solutions or suspensions of active compounds in sterile aqueous solutions, for example, aqueous propylene glycol or dextrose solutions. Such dosage forms can be suitably buffered, if desired.

Suitable pharmaceutical carriers include inert diluents or fillers, water and various organic solvents. The pharmaceutical compositions may, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. Thus for oral administration, tablets containing various excipients, such as citric acid may be employed together with various disintegrants such as starch, alginic acid and certain complex silicates and with binding agents such as sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tableting purposes. Solid compositions of a similar type may also be employed in soft and hard filled gelatin capsules. Preferred materials, therefor, include lactose or milk sugar and high molecular weight polyethylene glycols. When aqueous suspensions or elixirs are desired for oral administration the active compound therein may be combined with various sweetening or flavoring agents, coloring matters or dyes and, if desired, emulsifying agents or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin, or combinations thereof.

25 Methods of preparing various pharmaceutical compositions with a specific amount of active compound are known, or will be apparent, to those skilled in this art. For examples, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easter, Pa., 15th Edition (1975).

The following examples illustrate the preparation of the compounds of the invention. In the following examples, "Me" means methyl and "Et" means ethyl.

-20-


Example 14-(6-Chloro-2,3-dihydro-indol-1-yl)-7H-pyrrolo[2,3-d]pyrimidine

To 4-chloro-7H-pyrrolo[2,3-d]pyrimidine (10.0 g, 0.065 mol) in dry pyridine (90 ml) was added 6-chloro-2,3-dihydroindol-1-yl amine (14.8 g, 0.078 mol), and the mixture was heated in an 85°C oil bath for 2 days. The reaction was cooled to ambient temperature and concentrated in vacuo. The resulting residue was purified by flash chromatography on silica gel (375 g, 40 mm mesh) using 5% MeOH/CH₂Cl₂ to afford the title compound as a pink-orange solid (1.1 g, 4.3%): HRMS: Calculated 271.0750, Found 271.0729; anal. RP18-HPLC RT: 4.88 minutes.

The above compound was dissolved in minimal methanol and a solution of HCl in (HCl(g) bubbled into 2 ml Et₂O) was added dropwise until the mixture remained cloudy. The precipitated HCl salt was dried in vacuo, washed once with Et₂O, and dried in vacuo to constant mass (MP: 266°C (dec)).

Examples 2-3

The compounds of Examples 2 and 3 were made according to the method of Example 1 from 4-chloro-7H-pyrrolo[2,3-d]pyrimidine and the appropriate amine starting material.



Example no.	R	% Yield	RP18-HPLC RT	LC/MS (M ⁺)
2	6-Methyl	23	4.62	251
3	6-Chloro-5-fluoro	23	4.66	289

Example 41-(4-m-Tolylamino-pyrrolo[2,3-d]pyrimidin-7-yl)-ethanone

Following the procedure described in Example 1, (7H-Pyrrolo[2,3-d]pyrimidin-4-yl)-m-tolyl-amine was prepared from 4-chloro-7H-pyrrolo[2,3-d]pyrimidine and *m*-toluidine (34%): HRMS: Calculated 225.1140, Found 225.1131; anal. RP18-HPLC RT: 3.45 min; HCl salt MP: 219°C.

To (3-methyl-phenyl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine (0.168 g, 0.75 mmol) dissolved in hot acetonitrile (7 ml) was added sodium hydride (36 mg, 0.90 mmol, 60% dispersion in mineral oil). After stirring at ambient temperature for 0.75 hour, acetyl chloride (0.11 ml, 1.5 mmol) was added and stirring continued for 48 hours. The mixture was concentrated in vacuo, triturated in hot ethyl acetate, and filtered. The filtrate was concentrated in vacuo to give an orange solid residue. The solid was triturated in CH₂Cl₂ and filtered to afford the title compound as a light yellow solid (0.11 g, 55%): LC-MS: 267 (MH⁺); anal. RP18-HPLC RT: 3.53 minutes.

15


Example 54-(6-Chloro-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine

To a suspension of 4-hydroxy-pyrido[3,4-d]pyrimidine (0.103 g, 0.70 mmol) in dry pyridine (2 ml) cooled in an ice-water bath was added dropwise trifluoroacetic anhydride (0.20 ml, 1.4 mmol). After stirring for 0.5 hour, 6-chloroindoline (0.10 g, 0.66 mmol) and pyridine (0.14 g, 1.81 mmol) in dry DMF (1.5 ml) was added dropwise. The cold bath was allowed to warm to ambient temperature and then heated at 70°C for 3 hours. The reaction was cooled to ambient temperature and then added to methylene chloride (150 mL). The organic layer was washed with saturated sodium carbonate and water, and then dried over sodium sulfate. The solvent was removed by rotary evaporation and the residue purified by column chromatography (silica gel, 9/2/1 - CH₂Cl₂/hexanes/methanol) to give a pale yellow residue (0.048g, 28%): MP: 194-6°C; LC/MS: 283 (MH⁺).

20
25

Examples 6-8

The compounds of examples 6-8 were made according to the method of Example 5 from 4-chloro-pyrido[3,4-d]pyrimidine and appropriate amine starting materials.



Example #	R	Yield	HPLC RT	LC/MS (M ⁺)
6	6-Bromo-5-fluoro	45	3.64	359
7	6-Chloro-5-fluoro			315
8	6-Iodo			389

Example 9(7-Benzenesulfonyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-(3-ethynyl-phenyl)-amine

To 4-chloro-7H-pyrrolo[2,3-d]pyrimidine (1.0 g, 0.0065 mol) in dry THF (10 ml) under nitrogen at -78°C was added dropwise via syringe over 15 minutes *n*-butyllithium (2.5 M in hexane; 2.88 ml, 0.0072 mol). The cooling bath was removed and the solution was stirred for 1 hour. The resulting pyrrolo anion precipitated as a very fine white solid in a cloudy colorless solution. After the suspension was re-cooled to -78°C, benzenesulfonyl chloride (1.26 g, 0.0072 mol) was added neat via syringe. The resulting yellow mixture was allowed to warm slowly to ambient temperature overnight. The grey-white suspension was poured into 2% aqueous sodium bicarbonate (50 mL) and extracted with diethyl ether (4 x 20 mL). The combined extracts were washed with water and dried (potassium carbonate) and evaporated to give a light amber oil which crystallized from diethyl ether. The product was collected by filtration to 1.4 g (74%) of white solid: LC-MS= 294 (MH⁺); anal. RP18-HPLC RT: 4.40 minutes

The above compound was dissolved in MeOH and *m*-aminophenyl acetylene (0.159 g, 0.0013 mol), and the mixture was heated in an 85°C oil bath for 2 days. The

-23-

reaction was cooled to ambient temperature and concentrated in vacuo. The residue was triturated with diethyl ether to produce a white solid (0.234 g, 92%); LC-MS: 375 (MH⁺); anal. RP18-HPLC RT: 3.48 minutes.

Example 10

5 4-(6-Chloro-2,3-dihydro-indol-1-yl)-5H-pyrrolo[3,2-d]pyrimidin-6-ol

To a solution of 4-(6-Chloro-2,3-dihydro-indol-1-yl)-5-amino-6-methylacetylpyrimidine (541 mg, 1.55 mmol) in 40 mL of ethanol was added 25 mol% of 10% palladium on carbon (125 mg) and 0.11 mL of 1N HCl (1.55 mmol). The mixture was hydrogenated for 3 hours at 3.4 atm (50 psi)(H₂). The reaction is filtered through
10 Celite® and concentrated *in vacuo*. The brown residue was slurried in methanol and the white solid was filtered off (279 mg, 63%); LC-MS: 287 (M⁺); anal. RP18-HPLC RT: 5.61 minutes; MP: 250°C (dec).

Example 11

15 (3-Ethynyl-phenyl)-[7-(2-morpholin-4-yl-ethyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine

To a solution of 184 mg (1.4 mmol) of 4-(2-hydroxyethyl)morpholine in 10 mL of toluene was added 276 mg (2.0 mmol) of anhydrous potassium carbonate and then 32 mg (1.3 mmol) of 97% sodium hydride. After 30 minutes, 343 mg (1.0 mmol) of sulfonylated 4-chloro-7H-pyrrolo[2,3-d]pyrimidine was added and the reaction was
20 heated at 100°C for 2 hours. The reaction was then partitioned between ethyl acetate and water and the aqueous layer was extracted with two additional portions of ethyl acetate. The combined organic phase was washed with water, dried over magnesium sulfate and concentrated in vacuo. The residue was chromatographed on silica gel using 10% methanol/methylene chloride to give an amber oil (140 mg, 55%); LC-MS
25 267 (M⁺).

The above compound was dissolved in MeOH and *m*-aminophenyl acetylene (0.123 g, 0.001 mol), and the mixture was heated in a sealed tube in a 120°C oil bath for 12 hours. The reaction was cooled to ambient temperature and concentrated in vacuo. The residue was triturated with diethyl ether to produce a white solid (0.135 g,
30 74%); LC-MS= 348 (MH⁺); anal. RP18-HPLC RT: 3.33 minutes.

Example 12(3-Ethynyl-phenyl)-[7-(2-methoxy-ethyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine

Following a procedure analogous to that described in Example 11, the title product was prepared in 81% yield from 4-chloro-7-(2-methoxy-ethyl)-7H-pyrrolo[2,3-d]pyrimidine (1.0 eq) and *m*-aminophenyl acetylene (1.2 eq) in methanol: MP: 240-241°C; LC-MS: 292 (MH⁺); anal. RP18-HPLC RT: 4.16 minutes.

Example 13(3-Ethynyl-phenyl)-[7-[2-(2-methoxy-ethoxy)-ethyl]-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine

Following a procedure analogous to that described in Example 11, the title product was prepared in 81% yield from 4-chloro-7-[2-(2-methoxy-ethoxy)-ethyl]-7H-pyrrolo[2,3-d]pyrimidine (1.0 eq) and *m*-aminophenyl acetylene (1.2 eq) in methanol: MP: 240-241°C; LC-MS: 336 (M⁺); anal. RP18-HPLC RT: 4.29 minutes.

Example 14(7-Allyl-pyrrolo[2,3-d]pyrimidin-4-yl)-(3-ethynyl-phenyl)-amine

To 4-chloro-7H-pyrrolo[2,3-d]pyrimidine (1.3 g, 8.5 mmol) in dry THF (30 ml) was added sodium hydride (1.0 g, 0.25 mmol, 60% dispersion in mineral oil). After stirring at ambient temperature for 1 hour, allyl iodide (0.93 ml, 10 mmol) was added and stirring continued for 48 hours. The mixture was concentrated in vacuo, triturated in hot ethyl acetate, and filtered. The filtrate was concentrated in vacuo to give an orange solid residue. The solid was triturated in CH₂Cl₂ and filtered to afford 4-chloro-7-allyl-pyrrolo[2,3-d]pyrimidine as a light yellow powder (0.58 g, 36%): TS-MS: 194 (MH⁺).

To 4-chloro-7-allyl-pyrrolo[2,3-d]pyrimidine (0.5 g, 2.6 mmol) in dry MeOH (5 ml) was added *m*-aminophenyl acetylene (0.36 g, 3.1 mmol). The suspension was heated in a sealed pressure tube at 125°C for 20 hours. The reaction was cooled to ambient temperature and concentrated in vacuo. The resulting oil was purified by flash chromatography on silica gel (50 g, 40 mm mesh) using 3% MeOH/CH₂Cl₂ to afford the title compound as a yellow powder (0.29 g, 41%): TS-MS: 275 (MH⁺); anal. RP18-HPLC RT: 4.62 minutes.

-25-

Example 15N-(5-Iodo-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-N-m-tolyl-acetamide

To (3-methyl-phenyl)-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine (0.75 g, 3.4 mmol) dissolved in hot acetonitrile (30 ml) was added sodium hydride (0.16 g, 4.0 mmol, 60% dispersion in mineral oil). After stirring at ambient temperature for 0.75 hour, acetyl chloride (0.48 ml, 6.7 mmol) was added and the mixture was stirred for 48 hours. The mixture was concentrated in vacuo, triturated in hot ethyl acetate, and filtered. The filtrate was concentrated in vacuo to give an orange solid residue. The solid was purified by flash chromatography on silica gel (13 g, 40 mm mesh) using 1:3 ethyl acetate/hexanes to afford the title product as a yellow solid (0.21 g): TS-MS: 309 (MH⁺).

To 1-(4-m-tolylamino-pyrrolo[2,3-d]pyrimidin-7-yl)-ethanone (0.21 g, 0.79 mmol) in dry CH₂Cl₂ (5 ml) and dry MeOH (2 ml) was added Na₂CO₃ (0.17 g, 1.6 mmol). After stirring at ambient temperature for 0.75 hour, N-iodosuccinimide (0.35 g, 1.6 mmol) was added. The mixture was stirred at ambient temperature for 48 hours then concentrated in vacuo. The residue was diluted with CH₂Cl₂ and H₂O. The H₂O phase was extracted once with CH₂Cl₂. The organic phase was washed twice with H₂O, dried over Na₂SO₄, and concentrated in vacuo. The resulting residue was purified by flash chromatography on silica gel (11 g, 40 mm mesh) using 2% MeOH/CH₂Cl₂ to afford the title compound as a yellow solid (30 mg): TS-MS: 393 (MH⁺); anal. RP18-HPLC RT: 3.42 minutes.

20

Example 164-(6-Chloro-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine

6-Methyl-pyrido[3,4-d]pyrimidin-4-one (200 mg, 1.24 mmol), polymer-supported triphenylphosphine (2.06 g of 3.0 mmol P/g resin, 6.20 mmol) and anhydrous carbon tetrachloride (1.20 mL, 12.40 mmol) were combined in 1,2-dichloroethane (6 mL). The mixture was heated to 60°C under an atmosphere of dry N₂(g) for 18 hours. 6-Chloroindoline (1.1 eq.) was added and heating was continued at 60°C for another 18 hours. The triphenylphosphine-bearing resin was filtered off and washed several times with chloroform. The filtrate and washings were concentrated *in vacuo* and preparative reversed-phase (C18) chromatography utilizing a gradient of 15% to 70% MeCN/pH4.5, 50 mM NH₄OAc followed by lyophilization of the appropriate fractions afforded the title product (30%) as its free-base: MP: 232-234°C; LC-MS: 297 (MH⁺); anal. RP-HPLC: 4.33 minutes.


30

Examples 17-19

The compounds of examples 17-19 were made according to the method of Example 16 from 4-chloro-6-methyl-pyrido[3,4-d]pyrimidine and the appropriate amine starting materials.

5

10



15

Example no.	R	Yield	RP18-HPLC RT	LC/MS (M ⁺)
17	6-Bromo-5-fluoro	45	3.64	359
18	6-Chloro-5-fluoro			315
19	6-Iodo			389

Example 20

20

4-(4-Bromo-7-methyl-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine hydrochloride salt

The title product was prepared from 6-methyl-pyrido[3,4-d]pyrimidin-4-one (1.0 eq.) and 4-bromo-7-methyl-indoline (1.5 eq.) according to the procedure described for Example 16. The crude product from the filtrate was flash chromatographed on silica using EtOAc/hexanes/MeOH (9:2:1) to afford the free base which was converted to the hydrochloride salt as described for Example 23 (33% yield); MP: 232-244°C; LC-MS: 355, 357 (MH⁺); anal. RP-HPLC: 5.20 minutes.

25

Example 21

4-(6-Bromo-7-methyl-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine hydrochloride salt

30

The title product was prepared from 6-methyl-pyrido[3,4-d]pyrimidin-4-one (1.0 eq.) and 6-bromo-7-methyl-indoline (1.5 eq.) according to the procedure described for Example 16. The crude product from the filtrate was flash chromatographed on silica

-27-

using EtOAc/hexanes/MeOH (9:2:1) to afford the free base which was converted to the hydrochloride salt as described for Example 23 (34% yield); MP 212-229°C; LC-MS: 355, 357 (MH⁺); anal. RP-HPLC: 4.90 minutes.

Example 22

5 4-(6-Bromo-5-fluoro-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine hydrochloride salt

The title product was prepared from 6-methyl-pyrido[3,4-d]pyrimidin-4-one (1.0 eq.) and 6-bromo-5-fluoro-indoline (1.5 eq.) according to the procedure described for Example 16. The crude product from the filtrate was flash chromatographed on silica
10 using 2% MeOH / 98% CH₂Cl₂ to afford the free base which was converted to the hydrochloride salt as described for Example 23 (36% yield); MP: 262-264°C; LC-MS: 359, 361 (MH⁺); anal. RP-HPLC: 4.83 minutes.

Example 23

4-(6,7-Dimethyl-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine

15 To 4-chloro-pyrido[3,4-d]pyrimidine (200 mg, 1.21 mmol) in isopropanol (3 mL) was added 6,7-dimethylindoline (211 mg, 1.44 mmol) and pyridine (190 mg, 2.41 mmol). The mixture was heated to reflux under an atmosphere of dry N₂(g) for 6 hours. Solvent was removed *in vacuo* and the residue was dissolved in CHCl₃ and washed with saturated aqueous Na₂CO₃. The organic phase was dried over Na₂SO₄,
20 concentrated *in vacuo*, and flash chromatographed on silica in 45% acetone/hexanes to afford 60 mg of 4-(6,7-dimethyl-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine (LC-MS: 278 (MH⁺). This material was dissolved in a minimum volume of 10% MeOH in CH₂Cl₂, and 1 mole equivalent of HCl in diethyl ether was added dropwise with stirring. The mixture was diluted with diethyl ether (4 volumes) and the precipitated HCl salt of
25 the product was filtered and dried *in vacuo* (58 mg); MP: 248°C; GC-MS: 277 (M⁺); anal. RP-HPLC: 4.06 minutes.

Example 24

(3-Ethynyl-phenyl)-pyrido[3,4-d]pyrimidin-4-yl-amine

To 4-chloro-pyrido[3,4-d]pyrimidine (250 mg, 1.50 mmol) in N-methylpyrrolidin-2-one (0.5 mL) was added 3-ethynylaniline (212 mg, 1.81 mmol) and pyridine (237 mg, 3.0 mmol). The mixture was heated to 80°C under an atmosphere of dry N₂(g) for 3 hours. The reaction mixture was dissolved in CHCl₃ and washed with saturated aqueous Na₂CO₃, and brine. The organic phase was dried over Na₂SO₄, concentrated
30

-28-

in vacuo, and flash chromatographed on silica with a gradient of 40% to 70% acetone/hexanes to afford 120 mg of product. This material was precipitated as its hydrochloride salt by dissolution in minimal CHCl_3 , titration with HCl (1 eq.) in diethyl ether, and dilution with diethyl ether. The yellow salt was filtered and dried *in vacuo*
5 (133 mg): MP 233-235°C; LC-MS: 247 (MH^+); anal. RP-HPLC: 3.45 minutes.

Example 25

Benzo[b]thiophen-5-yl-pyrido[3,4-d]pyrimidin-4-yl-amine

To 4-chloro-pyrido[3,4-d]pyrimidine (250 mg, 1.50 mmol) in N-methylpyrrolidin-2-one (0.5 mL) was added benzo[b]thiophen-5-yl-amine (270 mg, 1.81 mmol) and
10 pyridine (237 mg, 3.0 mmol). The mixture was heated to 80°C under an atmosphere of dry $\text{N}_2(\text{g})$ for 3 hours. The reaction mixture was dissolved in CHCl_3 and washed with saturated aqueous Na_2CO_3 , and brine. The organic phase was dried over $\text{Na}_2\text{SO}_4(\text{s})$, concentrated *in vacuo*, and flash chromatographed on silica in 40% to 70% acetone/hexanes to afford 180 mg of product. This material was precipitated as its
15 hydrochloride salt by dissolution in minimal CHCl_3 , titration with HCl (1 eq.) in diethyl ether, and dilution with diethyl ether. The yellow salt was filtered and dried *in vacuo* (188 mg): MP: 280-282°C; LC-MS: 279 (MH^+); anal. RP-HPLC: 3.63 minutes.

Example 26

(3-Ethynyl-phenyl)-(5-p-tolyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

To a solution of 7-Benzenesulfonyl-4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidine (400 mg, 0.953 mmol) in 8 mL of toluene was added this (benzylidene acetone) dipalladium chloroform adduct (40 mg, 10% by weight), 4-methyl benzene boronic acid (260 mg, 1.91 mmol), sodium carbonate solution (2.86 ml of 1 M solution) and 3.0 ml of ethanol (absolute). The mixture was refluxed for seven hours. The reaction mixture
25 was filtered through Celite®, and the Celite® was washed with ethyl acetate. The filtrate was poured into a separatory funnel was washed with saturated sodium bicarbonate (100 mL, 2 times). The organic layer was dried over magnesium sulfate and concentrated *in vacuo*. The resulting green black oil was stirred in methanol (5 mL) and KOH (107 mg, 1.91 mmol in 3 mL of water) was added. After three hours, the
30 reaction mixture was concentrated and the residue dissolved in water. The solution was acidified with 1 N HCl and extracted with ethyl acetate (50 ml, 3 times). The organic layer was dried over MgSO_4 and concentrated *in vacuo*. The resulting oil was chromatographed on silica gel (180 g, 40 μm) using 30% ethyl acetate/hexanes to

-29-

provide the product as a white solid (225 mg 92%): TS-MS: 244, 246 (MH⁺). To 4-chloro-(5-p-tolyl)-7H-pyrrolo[2,3-d] pyrimidine (200 mg, 0.821 mmol) in dry methanol (4 mL) was added m-aminophenyl acetylene (106 mg, 0.903 mmol). The suspension was heated in a sealed tube at 120°C for 18 hours. The reaction was cooled to ambient temperature and concentrated *in vacuo* to a brown solid. The residue was chromatographed over silica gel (140 g, 40 μm) using 40% ethyl acetate/hexanes to provide the title compound as a light yellow solid (82 mg, 32%): MP: 245-247°C; anal. RP18-HPLC RT: 6.120 minutes.

Example 27

10 (3-Ethynyl-phenyl)-(5-thiophen-2-yl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

Following the procedure described in Example 26, the title compound was prepared in 40% yield from 7-Benzenesulfonyl-4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidine and thiophene-2-boronic acid followed by coupling with m-aminophenyl acetylene (27 mg, 0.233 mmoles) in methanol: MP: 230-232°C; TS-MS: 317 (MH⁺); anal. RP18-HPLC RT: 5.614 minutes.

Example 28

20 (3-Ethynyl-phenyl)-[5-(4-methoxy-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine

Following the procedure described in Example 26, 4-chloro-[5-(4-methoxy-phenyl)]-7H-pyrrolo[2,3-d] pyrimidine, the intermediate, was made in 74% yield from 7-benzenesulfonyl-4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidine and 4-methoxy benzene boronic acid: TS-MS: 260 (MH⁺). The title compound was synthesized by coupling the intermediate with m-aminophenyl acetylene in 75% yield: MP: 207-210°C; FAB MS: 341 (MH⁺); anal. RP18-HPLC RT= 5.640 minutes.

Example 29

25 (3-Ethynyl-phenyl)-[5-(3-nitro-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine

Following the procedure described in Example 26, 4-chloro-[5-(4-methoxy-phenyl)]-7H-pyrrolo[2,3-d] pyrimidine, the intermediate, was made in 74% yield from 7-benzenesulfonyl-4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidine and 3-nitro benzene boronic acid: TS-MS: 260 (MH⁺). The title compound was synthesized from coupling the intermediate with m-aminophenyl acetylene in 15% yield: MP: 189-190°C; TSMS: 469 (MH⁺); anal. RP18-HPLC RT= 4.67 minutes.

-30-

Example 30[5-(4-Chloro-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-(3-ethynyl-phenyl)-amine

Following the procedure described in Example 26, 4-chloro-[5-(4-methoxy-phenyl)]-7H-pyrrolo[2,3-d] pyrimidine, the intermediate, was made in 74% yield from 7-benzenesulfonyl-4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidine and 4-chloro benzene boronic acid: TS-MS: 260 (MH⁺). The title compound was synthesized by coupling the intermediate with m-aminophenyl acetylene in 33% yield: MP: 225-227°C; TSMS: 398 (MH⁺); anal. RP18-HPLC RT= 6.45 minutes.

Example 31

10 (3-Bromo-phenyl)-(6-bromo-5-phenyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine

Utilizing an analogous procedure as described in Example 26, 4-chloro-(5-phenyl)-7H-pyrrolo[2,3-d] pyrimidine, the intermediate, was made in 74% yield from 7-benzenesulfonyl-4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidine and 3-bromo benzene boronic acid: TS-MS: 230 (MH⁺). This second intermediate was coupled with 3-bromoaniline in methanol in a sealed tube to provide (3-bromo-phenyl)-[5-(4-chloro-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine in 75% yield: TSMS: 365 (MH⁺). (3-bromo-phenyl)-[5-(4-chloro-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine was brominated with N-bromosuccinimide impregnated onto silica gel. The reaction was complete after stirring at room temperature under nitrogen for 1.5 hours. The reaction was filtered and concentrated in vacuo to provide the title compound as a gray solid in 28 % yield (128 mg): M.P.: dec 300°C; TSMS: 445/447 (MH⁺/MH²⁺); anal. RP18-HPLC RT= 7.223 minutes.


-31-

CLAIMS

What is claimed is:

1. A compound of the formula


5




- 10 or a pharmaceutically acceptable salt thereof, wherein:

-32-


Z is a group of the formula




(Ia)




(Ib)



(Ic)




(Id)



(Ie)

or



(If)


15

wherein n is an integer from 0 to 2 and p is an integer from 0 to 3;


R¹ is H, C₁-C₆ alkyl or -C(O)(C₁-C₆ alkyl);

-33-

R^2 is phenyl or 1H-indazol-5-yl, wherein said groups are optionally substituted by 1 to 3 R^5 substituents, or R^2 is a group of the formula (Ii) or (Ij)



or



10


(Ii)

(Ij)

wherein p is an integer from 0 to 3 and n is an integer from 0 to 2;

or R^1 and R^2 are taken together to form a group of the formula (Ik)

15



20

;

(Ik)

25

wherein the dashed line indicates a single or double bond and m is an integer from 0 to 4;

each R^3 is independently H, $-C(O)OR^9$, or C_1-C_6 alkyl wherein said alkyl is optionally substituted by halo, $-OR^9$, $-NR^9R^{10}$, or $-C(O)OR^9$;

30

R^4 is R^3 , $-OR^9$, or $-NR^9R^{10}$;

each R^5 is independently halo, cyano, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, $-OR^9$, $-NR^9R^{10}$, nitro, or C_6-C_{10} aryl wherein said alkyl, alkenyl, alkynyl and aryl R^5

-34-

groups are optionally substituted by 1 to 3 substituents independently selected from halo, nitro, C₁-C₄ alkyl and -OR⁹;

R⁶ and R⁷ are independently H or R⁵;

R⁸ is H, -SO₂(C₆-C₁₀ aryl), -(CH₂)_q(5-10 membered heterocyclyl), C₂-C₆ alkenyl,
 5 C₁-C₆ alkyl, -(CH₂)_qO(CH₂)_q(C₁-C₆ alkoxy), -(CH₂)_q(C₁-C₆ alkoxy), -C(O)(C₁-C₆ alkoxy),
 or -SO₂(C₁-C₄ alkyl) wherein each q is independently an integer from 2 to 4;

each R⁹ and R¹⁰ is independently H or C₁-C₆ alkyl; and,

R¹¹ is trifluoromethyl, halo, nitro, -OR⁹, -NR⁹R¹⁰, cyano, C₁-C₄ alkyl, -S(O)_xR⁹
 wherein x is an integer from 0 to 2, -C(O)OR⁹, -OC(O)(C₁-C₄ alkyl), -C(O)NR⁹R¹⁰,
 10 -NR⁹C(O)(C₁-C₄ alkyl), -C(O)NHCO₂(C₁-C₄ alkyl), -NHCH₂C(O)NR⁹R¹⁰, -NHC(O)(C₁-C₄
 alkoxy), -NHOC(O)(C₁-C₄ alkyl), -NR⁹OR¹⁰, anilino, pyrrolidinyl, piperidinyl, azido,
 guanidino, phenyl, -C(O)(C₁-C₆ alkyl), benzenesulfonyl, allyl, thiophenyl, morpholino,
 piperazinyl, 4-(C₁-C₄ alkyl)-piperazinyl, phenylthio, benzenesulphonamido, 2-
 oxopyrrolidin-1-yl, 2,5-dioxopyrrolidin-1-yl, phenoxy, benzoyloxy, benzoylamino,
 15 -(CH₂)_wO(CH₂)_vOR⁹, -O(CH₂)_wO(CH₂)_vOR⁹, -O(CH₂)_wC(O)OR⁹, -O(CH₂)_wC(O)NR⁹R¹⁰,
 -(CH₂)_wS(CH₂)_vOR⁹, -NH(CH₂)_vO(C₁-C₄ alkyl), -NH(CH₂)_w(C₆-C₁₀ aryl),
 -NHC(O)(CH₂)_w(C₁-C₄ alkoxy), or -O(CH₂)_w(C₆-C₁₀ aryl), wherein w is an integer from 1
 to 4 and v is an integer from 2 to 4, and wherein the alkyl, heterocyclic, and aryl
 moieties of the foregoing R¹¹ groups are optionally substituted by 1 or 2 substituents
 20 independently selected from the group consisting of halo, C₁-C₄ alkyl, -OR⁹, -NR⁹R¹⁰,
 -C(O)OR⁹, -OC(O)(C₁-C₄ alkyl), -C(O)NR⁹R¹⁰, -NHC(O)(C₁-C₄ alkyl), nitro, imidazolyl,
 piperidino, morpholino, and piperazinyl;

with the proviso that where Z is a group of the formula (Ie), and R² is phenyl,
 then said phenyl is substituted by 1 to 3 substituents independently selected from C₁-C₆
 25 alkyl, C₂-C₆ alkynyl and halo, and one of R⁶ and R⁷ is halo or H and the other is as
 defined above; and,

with the further proviso that where Z is a group of the formula (Ia), (Ib), (Ic) or
 (Id), and R² is phenyl, then said phenyl is substituted by C₂-C₆ alkynyl.

2. The compound of claim 1 wherein R² is phenyl optionally substituted by
 30 1 to 3 R⁵ substituents.

3. The compound of claim 2 wherein R² is phenyl substituted by C₂-C₆
 alkynyl.

4. The compound of claim 3 wherein said alkynyl is ethynyl.

5. The compound of claim 1 wherein Z is a pyrrolo moiety of formula (Ie) or (If).

6. The compound of claim 1 wherein R¹ and R² are taken together to form an indole or indoline moiety of formula (Ik).

- 5 7. A compound selected from the group consisting of:
- 4-(6-Chloro-2,3-dihydro-indol-1-yl)-7H-pyrrolo[2,3-d]pyrimidine;
- 4-(6-Methyl-2,3-dihydro-indol-1-yl)-7H-pyrrolo[2,3-d]pyrimidine;
- 4-(6-Chloro-5-fluoro-2,3-dihydro-indol-1-yl)-7H-pyrrolo[2,3-d]pyrimidine;
- 1-(4-m-Tolylamino-pyrrolo[2,3-d]pyrimidin-7-yl)-ethanone;
- 10 4-(6-Chloro-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- 4-(6-Bromo-5-chloro-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- 4-(6-Fluoro-5-chloro-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- 4-(6-Iodo-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- (7-Benzenesulfonyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-(3-ethynyl-phenyl)-amine;
- 15 4-(6-Chloro-2,3-dihydro-indol-1-yl)-5H-pyrrolo[3,2-d]pyrimidin-6-ol;
- (3-Ethynyl-phenyl)-[7-(2-morpholin-4-yl-ethyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine;
- (3-Ethynyl-phenyl)-[7-(2-methoxy-ethyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine;
- (3-Ethynyl-phenyl)-[7-[2-(2-methoxy-ethoxy)-ethyl]-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine;
- 20 yl)-amine;
- (7-Allyl-pyrrolo[2,3-d]pyrimidin-4-yl)-(3-ethynyl-phenyl)-amine;
- N-(5-Iodo-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-N-m-tolyl-acetamide;
- 4-(6-Chloro-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6-Bromo-5-fluoro-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 25 4-(6-Chloro-5-fluoro-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6-Iodo-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(4-Bromo-7-methyl-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6-Bromo-7-methyl-2,3-dihydro-indol-1-yl)-6-methyl-pyrido[3,4-d]pyrimidine;
- 4-(6,7-Dimethyl-2,3-dihydro-indol-1-yl)-pyrido[3,4-d]pyrimidine;
- 30 (3-Ethynyl-phenyl)-pyrido[3,4-d]pyrimidin-4-yl-amine;
- Benzo[b]thiophen-5-yl-pyrido[3,4-d]pyrimidin-4-yl-amine;
- (3-Ethynyl-phenyl)-(5-p-tolyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;
- (3-Ethynyl-phenyl)-(5-thiophen-2-yl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;

(3-Ethynyl-phenyl)-[5-(4-methoxy-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine;
(3-Ethynyl-phenyl)-[5-(3-nitro-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-amine;
[5-(4-Chloro-phenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-(3-ethynyl-phenyl)-amine;
(3-Bromo-phenyl)-(6-bromo-5-phenyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amine;

5 and the pharmaceutically acceptable salts of the foregoing compounds.

8. A pharmaceutical composition for the treatment of a hyperproliferative disorder in a mammal which comprises a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

9. The pharmaceutical composition of claim 8 wherein said
10 hyperproliferative disorder is cancer.

10. The pharmaceutical composition of claim 9 wherein said cancer is brain, lung, kidney, renal, ovarian, squamous cell, bladder, gastric, pancreatic, breast, head, neck, oesophageal, gynecological, prostate, colorectal or thyroid cancer.

11. The pharmaceutical composition of claim 8 wherein said
15 hyperproliferative disorder is noncancerous.

12. The pharmaceutical composition of claim 11 wherein said disorder is a benign hyperplasia of the skin or prostate.

13. A pharmaceutical composition for the treatment of a hyperproliferative disorder in a mammal which comprises a therapeutically effective amount of a
20 compound of claim 1 in combination with an anti-tumor agent selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, and anti-androgens, and a pharmaceutically acceptable carrier.

14. A pharmaceutical composition for the treatment of pancreatitis or kidney
25 disease in a mammal which comprises a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

15. A pharmaceutical composition for the blastocyte implantation in a mammal which comprises a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

30 16. A pharmaceutical composition for treating a disease related to vasculogenesis or angiogenesis in a mammal which comprises a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

17. The pharmaceutical composition of claim 16 wherein said disease is

-37-

selected from the group consisting of diabetes, diabetic retinopathy, hemangioma, glioma, melanoma, Kaposi's sarcoma, ovarian cancer, breast cancer, lung cancer, pancreatic cancer, prostate cancer, colon cancer and epidermoid cancer.

18. A method of treating a hyperproliferative disorder in a mammal which
5 comprises administering to said mammal a therapeutically effective amount of a compound of claim 1.

19. The method of claim 17 wherein said hyperproliferative disorder is cancer.

20. The method of claim 19 wherein said cancer is brain, lung, squamous
10 cell, renal, kidney, ovarian, bladder, gastric, pancreatic, breast, head, neck, oesophageal, prostate, colorectal, gynecological or thyroid cancer.

21. The method of claim 18 wherein said hyperproliferative disorder is noncancerous.

22. The method of claim 21 wherein said disorder is a benign hyperplasia
15 of the skin or prostate.

23. A method for the treatment of a hyperproliferative disorder in a mammal which comprises administering to said mammal a therapeutically effective amount of a compound of claim 1 in combination with an anti-tumor agent selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating
20 antibiotics, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, and anti-androgens.

24. A method of treating pancreatitis or kidney disease in a mammal which comprises administering to said mammal a therapeutically effective amount of a compound of claim 1.

25. A method of preventing blastocyte implantation in a mammal which
25 comprises administering to said mammal a therapeutically effective amount of a compound of claim 1.

26. A method for treating a disease related to vasculogenesis or angiogenesis in a mammal which comprises administering to said mammal a
30 therapeutically effective amount of a compound of claim 1.

27. The method of claim 26 wherein said disease is selected from the group consisting of diabetes, diabetic retinopathy, hemangioma, glioma, melanoma, Kaposi's

-38-

sarcoma, ovarian cancer, breast cancer, lung cancer, pancreatic cancer, prostate cancer, colon cancer and epidermoid cancer.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 97/01393

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D471/04 C07D487/04 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	WO 97 02266 A (CIBA-GEIGY AG) 23 January 1997 see page 1 - page 2 ---	1,8
P,A	WO 96 40142 A (PFIZER INC.) 19 December 1996 cited in the application see the whole document ---	1,8
A	WO 96 31510 A (CIBA-GEIGY AG) 10 October 1996 see page 1 - page 5 ---	1,8
A	WO 95 23141 A (PFIZER INC.) 31 August 1995 cited in the application see page 73 - page 76 see page 84 ---	1,8
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

13 February 1998

Date of mailing of the international search report

10.03.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 97/01393

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 19774 A (WARNER LAMBERT COMPANY) 27 July 1995 see page 161 - page 164, line 14 see page 168, line 14 - page 171 ---	1,8
A	WO 95 15758 A (RHONE-POULENC RORER PHARMACEUTICALS INC.) 15 June 1995 see page 27 - page 28, line 14 ---	1,8
A	WO 94 13676 A (PFIZER INC) 23 June 1994 see page 1 - page 4 -----	1,8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 97/01393

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 18-20
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 18-20
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter: nal Application No

PCT/IB 97/01393

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9702266 A	23-01-97	AU 6414896 A	05-02-97

WO 9640142 A	19-12-96	AU 5479196 A	19-12-96
		CN 1141298 A	29-01-97
		CZ 9601641 A	11-12-96
		NO 962386 A	09-12-96
		PL 314641 A	09-12-96
		SI 9600184 A	30-04-97
		SK 72996 A	09-04-97

WO 9631510 A	10-10-96	AU 5273396 A	23-10-96
		CA 2214086 A	10-10-96
		EP 0819129 A	21-01-98

WO 9523141 A	31-08-95	AU 2972795 A	11-09-95
		CA 2183655 A	31-08-95
		CN 1141633 A	29-01-97
		CZ 9602413 A	16-07-97
		EP 0746554 A	11-12-96
		FI 963283 A	22-08-96
		HU 76291 A	28-07-97
		JP 9501953 T	25-02-97
		NO 963506 A	22-10-96
		PL 315941 A	09-12-96

WO 9519774 A	27-07-95	US 5654307 A	05-08-97
		AU 1731495 A	08-08-95
		AU 1833495 A	08-08-95
		BG 100614 A	31-03-97
		BG 100615 A	28-02-97
		CA 2177372 A	27-07-95
		CA 2177392 A	27-07-95
		CN 1139383 A	01-01-97
		CN 1139430 A	01-01-97
		CZ 9601970 A	17-09-97
		CZ 9601971 A	16-07-97
		EP 0742717 A	20-11-96
		EP 0741711 A	13-11-96
		FI 962855 A	13-09-96
		FI 962856 A	25-09-96

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 97/01393

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9519774 A		HR 950033 A	31-10-97
		HR 950034 A	31-10-97
		HU 74590 A	28-01-97
		HU 74589 A	28-01-97
		JP 9508126 T	19-08-97
		JP 9508127 T	19-08-97
		NO 963093 A	24-07-96
		NO 963094 A	24-07-96
		PL 315632 A	25-11-96
		PL 315633 A	25-11-96
		SK 89496 A	08-10-97
		SK 89596 A	06-08-97
		WO 9519970 A	27-07-95
		US 5679683 A	21-10-97
		ZA 9500441 A	10-10-95
		ZA 9500440 A	10-10-95
WO 9515758 A	15-06-95	US 5480883 A	02-01-96
		US 5710158 A	20-01-98
		AU 1305095 A	27-06-95
		US 5646153 A	08-07-97
		US 5714493 A	03-02-98
WO 9413676 A	23-06-94	AU 5666494 A	04-07-94
		CA 2150016 A	23-06-94
		CN 1097758 A	25-01-95
		CZ 9501584 A	17-01-96
		EP 0674641 A	04-10-95
		FI 935585 A	18-06-94
		HU 70505 A	30-10-95
		JP 7509726 T	26-10-95
		NO 952398 A	16-06-95
		NZ 258690 A	29-01-97
		PL 309357 A	02-10-95
		ZA 9309271 A	12-06-95